LA-UR-22-20420

Approved for public release; distribution is unlimited.

Title: 233U(n,g) measurements in the Unresolved and Fast Regime at LANSCE

Author(s): Couture, Aaron Joseph

Leal Cidoncha, Esther

Intended for: Tour discussion for NCSP program managers

Issued: 2022-01-18









Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher dientify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.





Delivering science and technology to protect our nation and promote world stability

- EST.1943 -

²³³U(n,γ) measurements in the Unresolved and Fast Regime at LANSCE


NCSP Site Visit

A. Couture and E. Leal Cidoncha

19 Jan 2022

Motivation


- Experimental ²³³U(n,γ) cross section data in the literature are scarce and were measured decades ago.
- New report [1] suggests that a simultaneous measurement with capture would be useful.
- For ²³³U fission is around one order of magnitude more likely than capture.
 - Good discrimination between gammas coming from capture and fission is required.
 - New measurement performed at LANL combining NEUANCE and DANCE.



 233 U(n, γ) and 233 U(n,f) cross sections from ENDF/B-VIII.


[1] M.T. Pigni, R. Capote and A. Trkov, Annals of Nuclear Energy 163 (2021) 108595.

A Brief Reminder on Time-of-Flight Measurements



Details vary with facility, but the basic principles are the same


Signatures of Capture: Why Detectors Matter




- Neutron Capture typically releases 6-10 MeV in gammaray energy
- Neutron energies do not significantly impact emitted energy

- Neutron capture measurements were often performed with C₆D₆ liquid scintillators
 - C₆D₆ has very low neutron sensitivity, but no gamma energy information
 - High Purity samples are always required
 - C₆D₆ has very low efficiency, typically requiring gram samples
- Calorimetric detectors can distinguish capture from decay based on total energy
 - High efficiency allows small samples
 - Isotopically mixed samples can be used if the isotopes have sufficiently different Q-Value
 - High segmentation offers gamma multiplicity information and limits individual crystal count rates

Greater than 2 order of magnitude improvement in S/N from DANCE cuts on Total Energy and Multiplicity



After Multiplicity and Total Energy Cuts



All data is from ²⁴⁰Pu measurement

²³³U Experiment at DANCE

- Uses the same fission:capture ratio technique used for ²³⁵U (Jandel et al.) and ²³⁹Pu (Mosby et al.)
- Used DANCE gamma calorimeter and NEUANCE fission tagger
- 20 mg sample of ²³³U provided by the National Isotope Development Center
- Ran in 2 beam cycles
 - Proposal was highest ranked proposal for DANCE in both cycles
 - Nov/Dec 2020
 - 21 days of beamtime
 - June/July 2021
 - 19 days of beamtime

DANCE and NEUANCE

DANCE (Detector for Advanced Neutron Capture Experiments)

 $4\pi BaF_2 \gamma$ -ray calorimeter composed by 160 crystals with an inner cavity of 17 cm radius [2].

Used to measure neutron capture cross section data on small quantities of radioactive isotopes.

We can measure E_n , E_{sum} , E_{γ} , and M_{γ} , providing more information than with C6D6 detectors.

NEUANCE (**NEU**tron detector array at d**ANCE**)

Neutron detector array that consists in 21 stilbene crystals arranged in a cylindrical geometry around the beam pipe [3].

Possibility to use a thick target.

Used to detect neutrons coming from fission and determine by coincidence with DANCE, the gammas coming from fission.

Those events are suppressed with a fission tag, and then the fission gamma shape is characterized with fission events to subtract the remaining fission background.


[2] M. Heil et al., Nucl. Instrum. Methods Phys. Res. A 459, 229 (2001).

[3] M. Jandel et al. Nuclear Inst. and Methods in Physics Research, A 882 (2018) 105-113.


NEUANCE

DANCE

Fission Tagging allows clear identification—and elimination—of fission gamma background



DANCE analysis cuts allow identification and *isolation* of different reaction channels





Evaluation collaboration and next analysis steps

- We met with Luiz Leal and Marco Pigni in early December (virtually)
 - Measurement technique, strengths, limitations, optimal neutron binning
 - Timeline for results (preliminary numbers to them in summer, final in fall)
 - Planned future discussions ~ quarterly including a meeting before the TPR
- Analysis
 - Optimize the fission tagging efficiency and acceptance
 - Extract capture yield
 - Determine capture cut efficiency
 - An independent determination of the fission and capture cross sections is possible, but would be a significant increase in the scope of work

The redesigned Lujan Target will enhance efficacy for keV cross sections

- Allow direct line-of-sight to W spallation target
- Improved intensity and resolution in the keV-MeV neutron energy range expand the range of measurements
 - Higher flux allows more flexibility for unstable isotopes
 - Improved resolution improves signal-to-noise in keV energy range





L. Zavorka, et al. NIMA 901 189 (2018)

²³³U(n,γ) Status and Prospects

- Capture measurements completed
 - Use of DANCE allows clean separation of capture, fission, and scattering channels
 - NEUANCE fission tagging with DANCE multiplicity gives reliable fission information
- Evaluation path and discussions are underway
 - Timeline for results (preliminary numbers to them in summer, final in fall)
 - Planned future discussions ~ quarterly including a meeting before the TPR
- Analysis is well underway to deliver a capture: fission ratio
 - An independent determination of the fission and capture cross sections is possible
- LANSCE neutron source improvements offer new measurement capability on small stable and radioactive samples





Delivering science and technology to protect our nation and promote world stability

- EST.1943 -